

James D. Kelly, II, M.D.
jkelly@calpacortho.com

Protocol for Reversed Total Shoulder Replacement

For Physical Therapist Use Only

General Guidelines:

- Maintain surgical motion early, but don't push it.
- Strengthen the surrounding musculature.
- Alternate between pool (if possible) and land therapy.
- Minimize heavy or excessive cyclic loads for the first 6 months.
- **No Manual Joint Mobilization!! This is a constrained prosthesis.**
- **NO Therapist Hands On Stretching... NONE!!! Allow patient stretching exercises.**
- **Most patients will only perform self directed therapy at home and then begin supervised PT at 6 weeks postoperative**

This is a gradual progression, not a stepped progression!!

Outpatient Rehab

Phase I: 0-3 weeks post op

- SLING:
 - o On except for:
 - Exercises
 - Dressing
 - Eating
 - Shower: Most have subcuticular closure with glue and/or water resistant dressing and may shower immediately.
- Pendulums and Codman's ex's
- Towel Slides or equivalent
- General conditioning (stationary bike, treadmill, etc)
- Maintain hand strength
- Maintain normal motion at the elbow/wrist

3-6 weeks post op.

- SLING
 - o OK to remove for eating, brushing teeth, shaving, reading, sitting in chair.

- Don't use arm to push up out of chair!!!
- Pool (if possible) for PROM.
- Supine AAROM (not to exceed surgical rom)
 - o FLEX, ABD, ADD, IR with towel, start ER at 6 weeks (minimize reps 5-10).
- Pool for PROM and AAROM (water is the assistance)
 - o Use combined motions and teach fluidity of movement.
 - o 10 reps with combined movements in pool.
- Light scapular strengthening (i.e. scapular setting, gentle MR scap protraction and retraction).
- Soft tissue mobilization to restore normal scapular mobility.
- Nerve glides as necessary

See pt 1-2x/week; mainly in the pool. Develop HEP for pt to work on PROM both in the pool and on land.

Phase II: 6-12 weeks

- Discontinue sling.
- It is fine to begin reaching behind back at this point.
- Continue with PROM.
- More AAROM on land
 - o Progress from SUPINE to SEATED, then to STANDING position.
- At **6 weeks** begin AROM on land, against gravity (straight planes only, no combined motions.)
- Pool – continue
 - o AAROM and AROM
- Can begin pulley work.
- Progress to some light closed-chain proprioceptive ex's (wall washing)
- Arm bike with no resistance.

See 1-2x/week, (increase frequency to 3x/week if ROM is not progressing.)

Phase III 12-18 weeks post op

- Generally, can begin progressing as tolerated.
- Can begin AROM with COMBINED MOTIONS at 3 months (on land, against gravity).
- Begin light strengthening
 - o Begin strengthening for lats, rhomboids, biceps/triceps, pecs, and deltoids.

James D. Kelly, II, M.D.
jkelly@calpacortho.com

- Pool – continue
 - o Use floats and paddles for resistive work in the pool.
- Increase scapular strengthening with theraband and light weights.
- Add more advanced proprioceptive ex specific for work, ADL's and sport.

See pt 2x/week, then transition to independent home ex program.

Please contact Dr. Kelly's Medical Assistant with any questions.

Olivia Stepp, Medical Assistant
Phone: (415) 668-8010 x 1020
Fax: (415) 928-1035
Email: ostepp@calpacortho.com